

Mannitol Salt Agar

Catalog : HB4128-11

Mannitol Salt Agar is used for the selective isolation and culture of Staphylococcus aureus.

Approximate Formula:

Ingredients	gm/liter
Pancreatic digest of casein	5.0
Peptic Digest of Animal Tissue	5.0
Beef Extract	1.0
D- Mannitol	10.0
Sodium Chloride	75.0
Phenol Red	0.025
Agar	15.0
Final pH7.4±0.2 at 25 °C	

*Adjusted and/or supplemented as required to meet performance criteria.

Directions:

Suspend 111.0g of the medium in one litre of distilled water or deionized water. Heat to boiling for 1 min with frequent agitation to completely dissolve the powder. Sterilize in an autoclave at 121 °C (15 lbs.) for 15 min.

Principle and Interpretation:

Mannitol Salt Agar is a nutritive medium due to its content of peptones and beef extract, which supply essential growth factors, such as nitrogen, carbon, sulfur and trace nutrients. The 7.5% concentration of sodium chloride results in the partial or complete inhibition of bacterial organisms other than staphylococci. Mannitol fermentation, as indicated by a change in the phenol red indicator, aids in the differentiation of staphylococcal species. Agar is a solidifying agent.

Appearance:

Dehydrated medium is a free-flowing yellowish-light red powder. The prepared medium is a kind of red transparent gel.

Precautions:

This medium is for laboratory use only. Dried medium which is past shelf life, caking or color variation cannot be used.

Storage conditions and Shelf life:

Mannitol Salt Agar must be stored tightly capped in the original container at 5-30°C. The dehydrated medium has a shelf life of 5 years from date of manufacturing. Prepared medium may be stored, out of direct light at 2-8°C.

Quality control:

Prepare the culture medium as per label directions. Inoculate and incubate at 36±1 °C for 24-48 hours.

Microorganism	Strains Number	Inoculum (CFU)	Growth	Recovery	Remarks
<i>Staphylococcus Aureus</i>	ATCC 6538	10-100	luxuriant	≥70%	Yellow colonies with yellow halo
<i>Staphylococcus epidermidis</i>	CMCC 26069	10-100	luxuriant	/	Red colonies
<i>Escherichia coli</i>	ATCC 25922	/	inhibited	/	/
<i>Staphylococcus Aureus</i>	ATCC 25923	10-100	luxuriant	≥70%	Yellow colonies with yellow halo
<i>Salmonella typhimurium</i>	ATCC 14028	/	inhibited	/	/

References:

1. United States Pharmacopeial Convention, Inc. 2008. The United States pharmacopeia 31/The national formulary 26, Supp. 1, 8-1-08, online. United States Pharmacopeial Convention, Inc., Rockville, Md.
2. European Directorate for the Quality of Medicines and Healthcare. 2008. The European pharmacopoeia, 6th ed., Supp. 1, 4-1-08, online. European Directorate for the Quality of Medicines and Healthcare, Council of Europe, 226 Avenue de Colmar BP907-, F-67029 Strasbourg Cedex 1, France.